NORESIDIE

Use Trifecta **CROP CONTROL**for treating and preventing problem mites, aphids, thrips, whiteflies, as well as botrytis, fusarium, and downy and powdery mildew.



ACTIVE INGREDIENTS: Thyme Oil 14%, Clove Oil 10%, Garlic Oil 9%, Peppermint Oil 4%, Corn Oil 3%, Geraniol 3%, Citric Acid 2%, Rosemary Oil 2% INERT INGREDIENTS: 53% Filtered Water, Soap, Isopropyl Alcohol, Vinegar Total 100%

Trifecta, LLC. - 516 West Central Avenue - South Williamsport, PA 17702 - 570-337-5125 - Trifecta.com.bz
Manufactured by Inventek Colloidal Cleaners, LLC.

THIS PRODUCT MAY BE USED ON CROPS INTENDED FOR HUMAN COMSUMPTION. THIS PRODUCT HAS NOT BEEN REGISTERED WITH THE UNITED STATES ENVIRONMENTAL PROTECTION AGENCY. TRIFECTÁ LLC REPRESENTS THAT THIS PRODUCT QUALIFIES FOR EXEMPTION FROM REGISTRATION LYNDER FIFRA LEGISLATION SECTION 250).



MOITLIA

STORE ONLY IN ORIGINAL CONTAINER IN A DRY PLACE INACCESSIBLE TO CHILDREN AND PETS. DO NOT REUSE EMPTY CONTAINERS. OFFER FOR RECYCLING IF AVAILABLE OR DISCARD IN TRASH.

AVOID CONTACT WITH EYES, SKIN OR CLOTHING, WEAR EY PROTECTION AND GLOVES WHEN HANDLING, IF IN EYE FLUSH WITH WITHER FOR AT LEAST 15 MINUTES, 15 ON SKIN WASH WITH PLENTY OF SOAP AND WATER. GET MEDICA ATTENTION IF IRRITATION PERSISTS. DO NOT APPLY THI PRODUCT INDILITED.

NOT INTENDED FOR AQUATIC USE.

HEALTHY GROWING IS HEALTHY LIVING. -SAFE FOR MOST CROPS -

NOREISE

DIRECTIONS FOR USE> Shake well before diluting.

FOR INFESTATIONS: For most applications dilute 2 oz. Crop Control per 1 gallon of water. For occasional severe infestations diluting up to 4 oz. Crop Control per 1 gallon of water may be necessary. Apply using an atomizer, misting system, or other spraying apparatus. Thoroughly wet plants until foliage is dripping including the underside of the leaves. **Crop Control should be applied** every other day up to 3 times per week until infestation is controlled. Apply Crop Control at the end of a light cycle or just before dusk to prevent foliar damage.

FOR PREVENTATIVE MAINTENANCE:

Dilute 1/2·1 oz. Crop Control per 1 gallon of water. Apply using an atomizer, misting system or other spraying apparatus. Thoroughly wet plants until foliage is dripping including the underside of the leaves. Preventative applications should be applied 1·2 times per week and preferably scheduled routinely on the same days/times each week. Apply Crop Control at the end of a light cycle or just before dusk to prevent foliar damage.



FLORIDA DISCLAIMER
FOR USE ON LOW-THC CANNABIS OR
MEDICAL CANNABIS BY APPROVED

NET CONTENTS: 32 FL. OZ. (946 ML)

KEEP OUT OF REACH OF CHILDREN