
Awakening

BIO-FERTILIZER





3 - 3 - 5

GUARANTEED ANALYSIS:

*Maximum Chlorine (Cl) 0.54%

3.00%
3.00%
5.00%

 Boron (B)
 0.10%

 Zinc (Zn)
 0.05%

 0.05% Water Soluble Zinc (Zn)

Derived from: Urea, Fish protein hydrolysate, Soy protein hydrolysate, Urea, Monopotassium phosphate, Potassium chloride. Boric acid. Zinc IDHA.

ALSO CONTAINS NON-PLANTFOOD INGREDIENT(S)

INERT INGREDIENTS: 95.00% 95.00% Total Other Ingredients (inert as non-plant food ingredients) Propolis (Bee attractant), Amino Acid Powder, Citric Acid, Glutamine, 5.00% Seaweed extract (Purpose, Microbe food) Disclaimer or Warranties. To the extent consistent with applicable law, GROPRO Inc. makes no otherwarranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of GROPRO Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein or to modify the warranties contained herein. To the extentconsistent with applicable law, GROPRO Inc. disclaims any liability whatsoever for special, incidental orconsequential damages resulting from the use or handling of this product.

Limitations of Liability: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at GROPRO Inc's election, the replacement of product

Contains Boron. Do not use on plants sensitive to Boron. Use of Boron on plants other than those recommended may result in serious injury to the plants.

PRECAUTIONARY STATEMENTS

Do not get in eyes, on skin or on clothing, Wash hands, forearms, and other exposed areas thoroughly after handling. Do not eat, drink or smoke when using this product. Do not breathe vapors, mist, or spray

//////// CAUTION ////////

KEEP OUT OF REACH OF CHILDREN



May be harmful if swallowed. Causes mild skin and eye irritation.

How can we help? 1-833-476-7761

2.5 Gal (9.46353 L)



MANUFACTURED BY

900 128th Street W, Burnsville MN 55337 Phone: 1-833-476-7761

www.groproag.com

·····5·-p·--9·--

9.81 lbs. per gallon @ 68 F FLORIDA FERTILIZER LICENSE NUMBER F003514 1 A, B, C, D, Root and Tuber Vegetables
Dosage per acre per application: 2-8 pints.
Two applications: 1) 100% sprouting, 2) 15 days after first treatment.

2 Leaves of Root and Tuber Vegetables
Dosage per acre per application: 2-8 pints.
Two applications: 1) 100% sprouting, 2) 15 days after first treatment.

3-07 A, B, Bulb Vegetable Group
Dosage per acre per application: 2-8 pints.,
Three applications: 1) At first flower. 2) 7-14 days later. 3) 7-14 days later.

6-22 A, B, C, D, E, F Legume Vegetable Group Dosage per acre per application: 3-8 pints. Two applications: 1) At 1st bloom. 2) At first pod.

8-10 A, B, C Fruiting Vegetable Group
Dosage per acre per application: 2-8 pints., Three applications: 1) At first
flower: 2) 7-14 days later. 3) 7-14 days later. Or during season at 2-4 pints per
application for clant growth

9 A, B Cucurbit Vegetables

Dosage per acre per application: 2-8 pints., Three applications: 1) At first flower. 2) 7-14 days later.3) 7-14 days later. Or during season at 2-4 pints per application for plant growth

10-10 A,B,C Citrus Fruit Group

Apply at a single shot of 3qts or 3 applications of 1 qt each per acres during the bloom cycle. Or during season at 2-4 pints per application for plant growth

11-10 Pome Fruit Group

Apply at a single shot of 3qts or 3 applications of 1 qt each per acres during the bloom cycle. Or during season at 2-4 pints per application for plant growth

12-12 A, B, C Stone Fruit Group

Apply at a single shot of 3qts or 3 applications of 1 qt each per acres during the bloom cycle. Or during season at 2-4 pints per application for plant growth

13-07 A, B, C, D, E, F, G, H, Berry and Small Fruit

Apply at a single shot of 1.5qts or 3 applications of 2 pints each per acres during the bloom cycle. Or during season at 2-4 pints per application for plant growth

14-12 Tree Nut Group

Apply at a single shot of 3-4qts or 3 applications of 1 qt each per acres during the bloom cycle. Or during season at 2-4 pints per application for plant growth

FIRST AID

Call a POISON CENTER or doctor immediately. • Rinse mouth. DO NOT induce vomiting. • Do not give anything by mouth to an unconscious person.
 Immediately call a POISON CENTER or doctor.

IF IN EYES:

 Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Call a poison control center or doctor for treatment advice.

F INHALED:

 Move person to fresh air and keep at rest in a position comfortable for breathing if they feel unwell.
 If not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-bornouth if possible.
 Call a POISON CENTER or doctor for treatment advice.

IF ON SKIN OR HAIR:

- Take off immediately all contaminated clothing. Rinse skin with water or shower. • Get medical attention if irritation develops or persists.
 Wash contaminated clothing before reuse.
- Tradit containmated electring percit of react.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact <u>I-800-424-9300</u> for emergency medical treatment information.